AgRISTARS

YM-N4-04456 JSC 1890**≅** 18908

A Joint Program for Agriculture and Resources Inventory Surveys Through Aerospace Remote Sensing

JANUARY 16, 1984

Yield Model Development

ARGENTINA CORN YIELD MODEL

(E84-10105) ARGENTINA CORN YIELD MODEL (National Oceanic and Atmospheric Administration) 15 p at A02/MF A01 CSCL 02C

N84-21923

Unclas G3/43 00105

SUSAN L. CALLIS CLARENCE SAKAMOTO



USDA/NOAA CIAD/MODELS BRANCH ROOM 200, FEDERAL BLDG 608 CHERRY STREET COLUMBIA, MO 65201











Lyndon B. Johnson Space Center Houston Texas 77058

ARGENTINA CORN YIELD MODEL

Ву

Susan L. Callis

and

Clarence Sakamoto

AISC Models Branch

January 16, 1984

	Report No. YM-N4-04456, JSC 18908	2. Government Acces	sion No.	3. Recipient's Catalog	g No.
4.	Title and Subtitle			5. Report Date January 16,	1984
	Argentina Corn Yield Model			6. Performing Organi	
7.	Author(s)			8. Performing Organia	ration Report No
	Susan L. Callis and Clarence	Sakamoto	ļ	10. Work Unit No.	
9.	Performing Organization Name and Address USDC/NOAA			TO. WORK ONLY 140.	
	CIAD/Models Branch Room 200, Federal Bldg, 608 (Cherry St.		11. Contract or Grant	No.
<u> </u>	Columbia, MO 65201			13. Type of Report ar	nd Period Covered
12.	Sponsoring Agency Name and Address National Aeronautics and Space Lyndon B. Johnson Space Center Houston, TX 77058		on _	14. Sponsoring Agency	Code
15	Supplementary Notes				
.5.	Sobbetueres à (40/62				
A model based on multiple regression was developed to estimate corn yields country of Argentina. A meteorological data set was obtained for the count averaging data for stations within the corn-growing area. Predictor variable for the model were derived from monthly total precipitation, average monthly temperature, and average monthly maximum temperature. A "trend variable" wincluded for the years 1965 to 1980 since an increasing trend in yields due technology was observed between these years.		for the country edictor variable everage monthly nd variable" was	y by es mean s		
17.	Key Words (Suggested by Author(s)) Multiple regression Predictor variables		18. Distribution Statement		
19.	Security Classif. (of this report)	20. Security Classif. (c	f this page)	21. No. of Pages	22. Price*
	Unclassified	Unclassifi	_	13	

INTRODUCTION

The purpose of this study was to select monthly weather variables that could be used to estimate yields of the Argentina corn crop. The corn-growing area in Argentina is indicated in Figure 1. Most of Argentina's corn is grown in a concentrated area in central Argentina where the climate is predominantly humid subtropical. The western edge of the soybean area in central Cordoba is semi-arid with warm to hot summers. There drought and high temperatures can be a problem during the growing season which begins with planting in October and November and extends to harvest in April through June.

METHOD

Multiple regression analysis of yield with selected agroclimatic indices was used to derive a suitable model. The regression equation for the corn model derived is:

$$\hat{Y} = \alpha + B_1T + B_2 (ETMETH_i) + B_3 (\Gamma X_i) + E$$
 where

 \hat{Y} = Estimated yield

 \propto = Constant

 B_j = Coefficients of the variables j = 1 - 3

T = Trend 1965-1980

ETMETH_i = ET - \widehat{ET} for month i: where \widehat{ET} = K·PET and K = $\overline{ET}/\overline{PET}$.

 TX_i = Maximum temperature for month i, and

E = Unexplained error.

Trend between the years 1965 and 1980 was determined from a plot of yields as shown in Figure 2. ET (evapotranspiration) minus $\widehat{\text{ET}}$ (climatically appropriate evapotranspiration) was used as an index representing soil moisture for plant growth. A more complete definition is given in the Appendix. Large positive $\widehat{\text{ET}}$ - $\widehat{\text{ET}}$ values suggest wet conditions.






Figure 1. DENSITY OF PLANTED AREA IN CORN 1977/78 CROP YEAR (Source: "Agronomic Characterization of the Argentina Indicator Region")





CORN PRODUCTION

CORN YIELDS

Figure 2. Plots of Production and Yield Versus Year for Argentina Corn.

In developing the models, various procedures of the Statistical Analysis System (SAS Institute, Inc., 1979) were used. The procedures used and the operations performed with each are summarized in the Appendix. The selected model had the highest \mathbb{R}^2 and included variables that were significant at the 10 per cent level and were agronomically meaningful.

DATA

Crop data for Argentina from 1960 to 1979 was obtained from the Foreign Agricultural Service (Alan Vandagrith, personal communication, 1982). A data set was created with year of yield as year of planting. Since the growing season spans two numerical years, the crop data was expressed in terms of "year + 1" (or harvested year) and the meteorological data corresponding to any yield included data for that year and "lagged" data for the previous year.

*/

The meteorological data was created using the general Argentina meteorological station file. This file was composed of data from several different sources, including the Monthly Climatic Data for the World and the Servicio Meteorological Nacional in Argentina (R.E. Jensen, C.M. Sakamoto, and S.E. Mummert; August, 1974). Stations inside the corn-producing area were averaged by province and weighted according to the percentage that their province contributed to the entire country's production. Figure 3 shows the corngrowing area in Argentina and associated stations; Table 1 lists the stations used and their weights.

The years between 1961 and 1980 (year of harvest) were used in the model since these years contained the most complete meteorological data.

PROCEDURES

Weather variables were selected from correlations and plots with yield.

These variables, along with TREND, were tried in regression equations. The

limited number of data years severely limits the number of variables that



Figure 3. Five major agricultural provinces in Argentina

STATION NAME	WE1 GH
BUENOS AIRES	.50
Pergamino Junin Nueve de Julio	
CORDOBA	.27
Rio Cuarto Bell Ville	
SANTA FE	.23
Rosario Casilda Las Delicias Parana	
Pergamino	

Table 1. Meteorological Stations Used for the Argentina Corn Model.

can be included in a regression model by limiting the number of degrees of freedom. This influenced the choice of model considered "best" for soybeans. Three weather variables plus TREND were considered the optimum model with 19 years of data. The three weather variables are:

ETMETH11......ET - ÉT for November

TX12......Maximum temperature for December

TX1......Maximum temperature for January.

The coefficient for the November variable was positive which reflects moisture requirements just after planting. The coefficients for maximum temperature for December and January were negative indicating that temperatures too high in the peak summer months during the tasselling and silking stages lead to stress on the plant. There is a critical growth period for corn. The statistics of the selected model are summarized in Table 2.

TEST RESULTS

A jackknife test was run on the model. In this test a year was eliminated from the crop data and the model was used to predict that year's yield. This process was done for each successive year beginning with 1961. The results were reasonable with the greatest difference (3.24 quintals) between predicted and actual yield in 1976. That year severe drought badly damaged the corn crop and harvest was hampered by rains. Test results are printed on Table 3 and plotted on Figure 4.

APPENDIX

Definition of Variables

The difference between ET and ET is an index of the amount of moisture available for plant growth. Soil moisture depletion is based on evapotranspiration (ET) estimates, determined as follows:

$$(ET)_{n} = \frac{(s)_{n-1}}{AWC} \left[\left\{ (PET)_{n} - (P)_{n} \right\} + (P)_{n} \right]$$
where

 $(ET)_n = "Actual" evapotranspiration,$

 $(S)_{n-1}$ = Available moisture at end of n-1 month,

AWC = Maximum water holding capacity,

 $(P)_n$ = Precipitation for month n, and

 $(PET)_n$ = Potential evapotranspiration for month n.

ET - ET measures the difference between the actual evapotranspiration and the "climatically appropriate" evapotranspiration, giving an indication of soil moisture supply and demand.

Statistical Analysis System Procedures Used

PROC CORR	Computes correlation coefficients between
	variables, including Pearson product-moment
	and weighted product-moment correlation

PROC STEPWISE FORWARD

Begins by finding the one-variable model that produces the highest
$$R^2$$
. For each of the other independent variables, FORWARD calculates F-statistics reflecting the contribution to the model if the variable were to be included.

PROC STEPWISE BACKWARD

Tr APP

Begins by calculating statistics for a model including all the independent variables. The variables are deleted from the model one by one until all the remaining variables produce F-statistics significant at the .10 level.

PROS STEPWISE STEPWISE

The stepwise method is a modification of the forward selection technique, differing in that variables already in the model do not necessarily stay there. After a variable is added (as in the forward selection method) the stepwise method looks at all the variables already included in the model and deletes any variable that does not produce an F-statistic significant at the .10 level. Only after this check is made and the necessary deletions accomplished can another variable be added to the model.

PROC STEPWISE MAXR

(Maximum R² improvement) Unlike the three techniques above, this method does not settle on a single method. Instead it looks for the "best" two-variable model, the "best" three variable model, and so forth.

PROC PETM

Uses latitude and mean monthly temperature to calculate Thornthwaite's potential evapotranspiration for each month.

PROC ZINDEX

Uses monthly PET's, pre ipitation, SS (beginning moisture in surface layer), AWCS (available water capacity in surface layer, SU (beginning moisture in the underlying layer), and AWCU (available water capacity in the underlying layer) to calculate Palmer's soil moisture budget, drought index Z, ET, and ET.

STATICAL AVALYSTS SYSTEM

THE PROPERTY OF THE PROPERTY O

r

IN:41 THESTAY. MAR

	. S V
	<i>J</i> .
	VERSION O
PPOPTED.	SUPPORTED
Ē	T.E
101 51	OHTATM
345	2
AN MING: THIS VIOSION OF SAS IS NOT SUBBORTED.	COMMITMS PERMESENTATIVE TO OMIAIN THE SUPPORTER VERSION OF SAS.
HI :Union	A SALL HOLD
	tons facili
	Collect

		HANGE 4A J ELLINES	HACKER J ELIMINATION PROCFOURE FOR DEPENDENT VAPIABLE YIELD	WIDENT VAPTABLE YTE	61.	
=	d in Avolved to bullere to	0 = 0.460 5 0	= 10,9 47494714 ((U) =	5.00000000	ļ	
		7)C	511+ OF SOUA-FS	MF AN SQUADE	•	PROBYE
	97.53193 F0090 T0T4L	*=*	151,29894507 20,46504393 172,16400000	37.8247190.5 2.04650439	14.13	0.0041
		JIITVA E	SIN turin	TYPE II SS	•	Panase
	[:te:/cf.bt :f:/: :f:/:	(A) A A A A A A A A A	0.1553a+R2	44.59797920 7.0084601	4.5 . C	0.000.0
	1212	-0.57.19.37.60	107/01/20 07/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00/01/20 00	8.91427490 17.69568470	\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\	0.065

TABLES OF THE MODEL AND STONIFICANT AT THE USING LEVELS

Table 2. Statistics of Argentina Corn Model.

JAS.

ークうならんてみりの1ろうなら

CO.WIES

F 10E

14:41

Results of Jackknife Test for Argentina Corn Model. Table 3.

ないると

1

Quintals

A 176 , 🛊

.4 -

figure 4. Argentina Corn Model.